

Technology & Engineering

May 8, 2008

US EPA Office of Pollution Prevention and Toxics
-EPA East Building Room 6428

Attn: Section 8(e)

1201 Constitution Avenue, NW

Washington, DC 20004

8 E H Q - 0 7 - 1 6 9 0 8

03 MY 12 AN 6: (

SUBJECT: TSCA 8(e) Notice

Dear TSCA Section 8(e) Coordinator:



On behalf of Akzo Nobel Polymer Chemicals, LLC we are submitting results of a Reproduction/Developmental study conducted with hexaneperoxoic acid, 2-ethyl-, 1,1-dimethylethyl ester (CAS# 3006-82-4).

The test material was prepared in corn oil which was also used as the vehicle control. Dose groups were: control, 100, 300, and 1000 mg/kg. Doses were administered daily by oral gavage to male rats for 41 days and to female rats during pre-mating, mating, gestation, and lactation through 3 days post partum.

All animals survived until necropsy and no abnormal macroscopic or microscopic findings were noted. Clinical signs of toxicity were noted in the high dose males and females and included moving the head through the bedding material after test article administration. During the last two days of gestation or the first two days of lactation, 7 and 4 high dose dams, respectively had ruffled fur and a generally bad condition. During lactation, the mean body weight gain of the dams in the high dose group was statistically significantly decreased. Testes weights in high dose males were not affected by treatment although the organ/body weight ratio was statistically significantly increased.

Mating performance and fertility, the corpora lutea count, the implantation rate, and gestation length were not influenced by treatment with the test material. The mean pre-implantation loss was increased in treated groups but was not dose-related. Pre-implantation loss was reported previously for the range finding portion of this study (8EHQ-07-16908).

Post-implantation loss was statistically significantly increased in the high dose group. The total implantation loss was 15, 13, 10 and 49 for the vehicle control, low, mid and high dose groups, respectively. One dam in the high dose group had total litter loss corresponding to 8 lost implantation sites.

The number of living pups at first litter check was reduced when compared with controls. The mean number of living pups was 13.5, 12.1, 11.5, and 9.5 for the control, low, mid and high dose groups, respectively. Number of dead pups in the control, mid and high dose groups were 3, 2, and 40, respectively.

Postnatal loss was statistically significantly increased in the high dose group and low dose group. This finding in the low dose group was considered incidental as only 1/10 dams was affected by cannibalization of the pups. In the control, low, mid and high dose groups postnatal loss was 3, 12, 0 and 23 dead pups noted in 1, 1, 0 and 5 litters, respectively.

Mean high dose pup weight was statistically significantly reduced up to day 4 post partum. No treatment related macroscopic findings were observed in pups.



CONTAINS NO CB

Akzo Nobel Inc. 525 West Van Buren Street Chicago, Illinois 60607-3835 Tel. (312) 554 7000 Fax (312) 544 7125 Please contact me at (312) 544-7061 if you have any questions regarding this letter.

Sincerely,

Louette Rausch

Senior Staff Toxicologist

Akzo Nobel Services Inc./T&E

Youth Rouseh

525 W. Van Buren

Chicago, II 60607

From: Origin ID: GYYA (312)544-7315

FRANK THÖRNE

AKZO NOBEL SERVICES INC 525 WEST VAN BUREN STREET

CHICAGO, IL 60607



CLS120787/21/24

SHIP TO: 2025468930

BILL SENDER

TSCA Section 8(e) Coordinator U.S. EPA - Offc Piltn Prvntn & Txcs 1201 Constitution Avenue, NW **EPA East Building, Room 6428** Washington, DC 200043302



Ship Date: 08MAY08 ActWgt: 1 LB

System#: 3312538/INET8011 Account#: S ********



Invoice # PO# Dept#

TRK#

0201

FRI - 09MAY 7989 3715 1920

A1 STANDARD OVERNIGHT

NE RDVA

20004 DC-US IAD



Shipping Label: Your shipment is complete

- 1. Use the 'Print' feature from your browser to send this page to your laser or inkjet printer.
- 2. Fold the printed page along the horizontal line.
- 3. Place label in shipping pouch and affix it to your shipment so that the barcode portion of the label can be read and scanned.

Warning: Use only the printed original label for shipping. Using a photocopy of this label for shipping purposes is fraudulent and could result in additional billing charges, along with the cancellation of your FedEx account number.

Use of this system constitutes your agreement to the service conditions in the current FedEx Service Guide, available on fedex.com. FedEx will not be responsible for any claim in excess of \$100 per package, whether the result of loss, damage, delay, non-delivery, misdelivery, or misinformation, unless you declare a higher value, pay an additional charge, document your actual loss and file a timely claim. Limitations found in the current FedEx Service Guide apply. Your right to recover from FedEx for any loss, including intrinsic value of the package, loss of sales, income interest, profit, attorney's fees, costs, and other forms of damage whether direct, incidental, consequential, or special is limited to the greater of \$100 or the authorized declared value. Recovery cannot exceed actual documented loss. Maximum for items of extraordinary value is \$500, e.g. jewelry, precious metals, negotiable instruments and other items listed in our Service Guide. Written claims must be filed within strict time limits, see current FedEx Service Guide